DOCUMENT RESUME

ED 189 198 TM 800 440

AUTHOR Blumberg, Phyllis; Felner, Joel
TITLE A Practical Methodology for the Systematic Development of Multiple Choice Tests.
INSTITUTION Illinois Univ., Urbana. Medical Center.
SPONS AGENCY National Heart and Lung Inst. (DHEW/PHS), Bethesda, Md.
PUB DATE Apr 80
CONTRACT NIH-1-HU-6-2969
NOTE 22p.; Paper presented at the Annual Meeting of the National Council on Measurement in Education (Boston, MA, April 8-10, 1980).

EDRS PRICE MF01/PC01 Plus Postage.
DESCRIPTORS Computer Assisted Testing; Course Content; Higher Education; *Item Analysis; Item Banks; Medical Education; *Multiple Choice Tests; *Test Construction; Test Items
IDENTIFIERS Facet Analysis; *Guttman Facet Design Technique; *Test Equivalence

ABSTRACT

Using Guttman's facet design analysis, four parallel forms of a multiple-choice test were developed. A mapping sentence, logically representing the universe of content of a basic cardiology course, specified the facets of the course and the semantic structural units linking them. The facets were: cognitive processes, disease priority, specific disease, and clinical judgment stage. Developing items from specification strings was simpler and quicker than developing items from objectives without specific guidelines. Due to logistical problems, only two of the four parallel forms originally planned were actually used. Although this increased the number of times each item was used, it did not violate the theoretical methodology, nor the notion that the tests were parallel. The tests will be field tested with fourth-year medical students at five medical schools, to establish their parallel forms reliability. (CP)

****
* Reproductions supplied by EDRS are the best that can be made *
* from the original document. *
****

A PRACTICAL METHODOLOGY FOR THE SYSTEMATIC DEVELOPMENT OF EQUIVALENT MULTIPLE-CHOICE TESTS

ED189198

U S DEPARTMENT OF HEALTH, EDUCATION & WELFARE
NATIONAL INSTITUTE OF EDUCATION

THIS DOCUMENT HAS BEEN REPRODUCED EXACTLY AS RECEIVED FROM THE PERSON OR ORGANIZATION ORIGINATING IT. POINTS OF VIEW OR OPINIONS STATED DO NOT NECESSARILY REPRESENT OFFICIAL NATIONAL INSTITUTE OF EDUCATION POSITION OR POLICY

Phyllis Blumberg, Ph.D.
University of Illinois at the Medical Center

and

Joel Felner, M.D.
Emory University School of Medicine

"PERMISSION TO REPRODUCE THIS MATERIAL HAS BEEN GRANTED BY
P. Blumberg
TO THE EDUCATIONAL RESOURCES INFORMATION CENTER (ERIC)."

Paper presented at the annual meeting of the National Council on Measurement in Education in Boston, Massachusetts, April, 1980.

This research was supported by The National Heart, Lung, and Blood Institute Public Health Service, Department of Health, Education and Welfare, NIH # NO 1-HU-6-2969.

TM 800440

The authors wish to acknowledge the assistance of the other project members: the cardiologists - Michael S. Gordon, Gordon Ewy, Alan Forkner, Ira Gessner, Joan Mayer, and Robert Waugh; and the other educational consultants - Christine McGuire, Mary E. Risley, and Abdul Sajid. The authors are especially indebted to John Engel who provided theoretical and technical assistance throughout all stages of the development of this paper.

A Practical Methodology for the Systematic Development of Equivalent Multiple-Choice Tests

Phyllis Blumberg, Ph.D.
University of Illinois at the Medical Center

and

Joel Felner, M.D.
Emory University School of Medicine

This paper discusses how parallel multiple-choice question tests were developed using Guttman's facet design analysis. Facet design analysis provides a logical link between the instructional content and the items because the items are specified by the facet analysis of the objectives. A mapping sentence was developed from the facet analysis which defined the universe to be tested. This mapping sentence was used to specify a test blueprint for the parallel exams and also specified the particular dimensions of each item. This paper shows how facet design analysis is a practical way to develop parallel exams.

A PRACTICAL METHODOLOGY FOR THE SYSTEMATIC DEVELOPMENT OF EQUIVALENT MULTIPLE-CHOICE TESTS

OBJECTIVES

The development of a detailed blueprint of an exam including the specifications of test content and the number of questions per content area is an important step in test development. When alternate forms of an examination are needed, test constructors most often use the process of developing a content-by-cognitive behavior matrix as a way of a priori insuring functionally equivalent tests. However, most test development reference books (e.g. Popham, 1975 and Thorndike, 1971) do not discuss rules for constructing these matrices or for selecting items. These matrices may be a product of the test developers concept structure and their own idiosyncracies. Often, sets of items have not been shown to be equivalent by any empirical method (Bormuth, 1970).

A number of new approaches have been proposed which attempt to define rules for generating items. These item-generated mechanisms provide for the construction of tests which are content valid in a logical as well as judgmental way (Millman, 1974 and Martuza, 1977). One of these new approaches, the facet design approach, is employed here to develop equivalent multiple-choice tests. The purpose of this paper is to illustrate a method of developing alternate forms of tests according to logical rules of equivalence of facet design analysis.

Theoretical Orientation

Facet design was developed by Guttman (1969) for the purpose of defining a research domain. Facet design specifies the limit of the domain as well as the ordering of all of the subcomponents of the domain. It is a powerful tool in defining a problem, structuring relationships within and among variables, and dealing with the problems of equivalence (Jordan, 1978). A system similar to a Cartesian coordinate system is used to specify the relationship of subcomponents of the domain. Instead of drawing the numerous coordinates on graph paper, a mapping sentence is constructed (Engel, 1975).

A mapping sentence provides an integrated system for defining the universe of relevant concepts. It is composed of two parts: the facets and the phrases linking the facets together. The facets are categories of relevant content. The phrases linking these categories suggest the semantic structural unity which integrate all of the facets together. Each facet is defined in terms of specific information which is presented in a list. A facet element is a particular example from the list of a relevant concept. Figure 1 shows a sample mapping sentence. Thus, a mapping sentence, through the use of facets and facet elements, analyzes the domain to be sampled into all of its relevant component parts (Millman, 1974, Martuza, 1977, and Shye, 1978).

Insert Figure 1 About Here

One specific element from each facet is combined to form a facet string. A facet string defines a particular segment of the universe. Theoretically, the number of possible facet strings is equal to the number of combinations of all the coordinates. However, some of these facet strings are null sets when real data are applied.

Millman (1974), Engel (1975), Martuza (1977), and Berk (1978) propose the use of facet-design analysis for test-item construction. Initially the test constructor conceptually analyzes the content to be tested by dividing the content into its component parts. The result of this analysis yields a mapping sentence of the universe. Specific items can be generated from facet string specifications.

After reviewing various techniques for test specifications and item-generation, Engel (1975) concluded that the facet-design approach is the only approach which is explicitly based on a meta-theory of content representativeness. This advantage of a theoretical base makes this approach fundamentally attractive since it is operational for constructing multiple-choice tests. Facet-design analysis provides a logical link between the instructional content and the multiple-choice items because the items are specified by the facet-design which follows from the objectives. Unless a theoretically based methodology such as facet design is used, there is no a priori way of justifying the necessary assertion that the items included in the parallel forms define the same kind of behaviors. By constructing test specifications and items on the basis of a facet analysis, it becomes possible, in principle, to devise parallel forms in terms of content.

Methodology

Task

A test specialist was asked to help cardiologists develop four parallel multiple-choice tests examining the content of a basic cardiology course as part of a larger research project. This project involved five medical schools. Each test was to be composed of about 100 items. First the cardiologists identified twenty-three diseases that were pertinent to the students' level of learning. Next, these diseases were ranked high, medium or low priority based on frequency of occurrence. The test specialist employed facet design analysis in order to insure content domain equivalence.

The test specialist studied the objectives and developed a mapping sentence which had four major facets and at least three elements within each facet. Figure 2 shows the mapping sentence used for the test specification and item development. The elements have been ordered on the following dimensions: Facet A - cognitive processes employed which are ordered according to an adaptation of Gagne's (1970) and Bloom's, et. al. (1956) cognitive hierarchies; Facet B - disease categories which are ranked from high to low priority in terms of the objectives; Facet C - specific diseases which are ordered alphabetically; and Facet D - the stages in clinical judgment which are ordered by the chronological sequence of clinical judgments.

Insert Figure 2 About Here

Theoretically, the next step would be to list all of the possible 2,592 facet strings. (There is a total of 39 different coordinates from four facets - 6 X 3 X 24 X 6.) This was an unrealistic step especially since many of the facet strings do not reflect reality. For example, the use of visual information such as x-rays or electrocardiograms only enter into selected steps such as data

gathering (D_1), interpret laboratory data (D_3) and diagnose patient problems (D_4). While computer techniques are available to list the strings, it was decided that in this particular application, it would be sufficient to allow the content specialists to select the salient strings.

Test Specification

One cardiologist determined the overall blueprint for the exam. For example, he specified that about one-third of the items should test the student's ability to describe and interpret chest x-rays, electrocardiograms, phonocardiograms and echocardiograms (e.g., A_2 and A_4). From this general specification of the types of desired questions, the particular facet strings were selected which fulfilled these general criteria.

Each disease was ranked high, medium or low priority in the objectives. The test specifications stated the approximate number of items for each disease for the three ranked priorities. However, the particular disease was not specified in the test blueprint. Thus, Facet C, the name of the disease was incorporated into Facet B in the test specification facet string.

Table 1 shows the test specification blueprint defined in facet strings. The three facets used in test specification are: A. cognitive processes required, B. priorized diseases and D. the step within the clinical process. The number after each Facet letter refers to the specific element of the Facet as identified on Figure 2. Specification $A_1B_2D_4$ means that items should pertain to, "Recall of factual information relating to the diagnosing of patients with high priority diseases." This specification was used to generate two items. In order to insure that the four tests would be parallel in terms of content, one set of facet strings determined the test specification blueprint for all of the examinations.

The visual information referred to in Facet A_2- verbally describe visual information and Facet A_4- interpret visual information includes the use of the

following diagnostic tests: ambulatory electrocardiographic monitoring (Holter monitoring), angiography cardiac catheterization, chest x-ray, cineangiogram-coronary-arteriography, echocardiogram, electrocardiogram, impulse tracing, phonocardiogram, radionuclide imaging (Mycardial-scanning), and treadmill stress test. The same diagnostic test used on one exam was used on the other three exams with that facet string. For example, since Exam A required students to interpret a chest x-ray during the data gathering phase from a patient with mitral stenosis, Exams B, C and D also required students to interpret chest x-rays during the data gathering phase from patients with high priority diseases.

Facets A, B, and D are used for test specification, whereas Facets A, C, and D are used for item specification. The rationale for the use of these facets is explained earlier.

Insert Table 1 About Here

Item Specification

It was assumed that at least one item could be written for each disease which fulfilled the criteria of a facet string in Table 1. For example, the facet string for the test specification "$A_1B_1D_1$" specified that the item should recall factual information relating to the data gathering step on a high priority disease. There were ten high priority diseases. Therefore, at least ten different items could be developed from this facet string since it was assumed that knowledge about the ten diseases are interchangeable. These ten items could be assigned to ten parallel examinations. Since only four parallel examinations were required, only four such items had to be developed. However, for the facet string "$A_1B_1D_1$" two items per examination were required because of the item specification developed (see Table 1). Thus, eight separate items had to be developed which required the recall of factual information pertaining to ordering laboratory tests for high

priority diseases. Preferably these eight items would each deal with a separate high priority disease.

As indicated in the above paragraph, three facets (A, C, and D) were used for the item specification. Since all of the diseases were assigned a priority ranking, the priority facet (B) was not necessary when the individual diseases were considered in the actual item specification. The test specification facet string $A_1B_1D_1$ resulted in the following ten item specification facet strings since there were ten high priority diseases:

$A_1C_2D_1$	$A_1C_{10}D_1$
$A_1C_3D_1$	$A_1C_{12}D_1$
$A_1C_5D_1$	$A_1C_{13}D_1$
$A_1C_7D_1$	$A_1C_{14}D_1$
$A_1C_8D_1$	$A_1C_{15}D_1$

At least one item could be developed from each item specification facet string.

Item Development

The cardiologists had already begun to develop a pool of items. Only those items which met the test specification facet string requirements were selected for the pool. In some cases several items were in existence. Instead of rejecting good items, these parallel items were placed in the pool.

After the classification of the existing items it became clear that more items were needed to meet the exam specifications. The cardiologists were then asked to develop items according to the particular test specification facet strings that were lacking the required number of items as shown in Table 1. The item pool contained sets of items grouped according to the test specification facet strings from which they were derived.

Test Development

Since the cardiologists did not have the time to develop four hundred items, they asked the test specialist to recommend a way of developing useful exams using fewer items. After a discussion with one cardiologist, it was decided that 50% of each exam could have common items. Next the test specialist assigned items to each exam based upon the test specification table of facet strings (Table 1). Items were assigned randomly to one of four exams except with the stipulation that if more than one item was required for that facet string per exam, each item in an exam from that facet string would be concerned with a different disease. A running tally was kept on the items as they were assigned to exams in order to insure that there would be the correct number of items for each high, medium, and low priority disease. Frequently items had to be juggled from one exam to another. Yet, no matter how the items were juggled, all of the test specification criteria (e.g., Table 1 and the frequency of items per disease) could not be met with the existing pool. For example, on one exam there were too many items on normal patients and not enough items on aortic stenosis. Therefore, a cardiologist was asked to develop more items to be placed in specific slots on each exam on the basis of item specification facet strings. An item specification for each of the four parallel exams was developed. Table 2 shows the item specification for Exam A.

Insert Table 2 About Here

Test Use

The four exams were pilot tested. During the review process, it was decided that only two exams would be used due to logistical problems and problems with the difficulty or wording of specific items. The items on these two exams were derived from the original four parallel exams.

These exams will be used for fourth-year medical students in a cardiology rotation at five colleges of medicine in the United States as part of a larger experimental, simulated instructional program. The performance of these students on these exams will be analyzed to test the hypothesis that these exams are parallel in terms of content based on the facet design analysis conducted. The Guttman-Lingo computer analysis as well as factor analysis will be used to test this hypothesis.

Discussion

Periodically, throughout the test development process, the steps defined by the facet design analysis were modified. These slight modifications did not sacrifice the rational approach to test development. Ideally, one should initially list all of the possible facet strings and eliminate the unrealistic ones. However, this is not a practical way of developing test blueprints since the number of dimensions is usually large. The test developers selected facet strings from the mapping sentence without actually listing all possibilities. Since the mapping sentence listed all of the relevant coordinates, the relevant facet strings were easy to visualize.

According to the rigorous interpretation of Guttman's theory, all of the items should be developed once the test blueprint defining the facet strings is specified (Martuza, 1977). Yet, realistically, this may not be the case. Teachers and test developers often rely on a pool of previously constructed and used items.

The test specification facet string used three facets; the item specification facet string used three facets, two of which were the same as in the test specification facet strings. An alternative approach would have been to develop two mapping sentences. The cardiologists believed that the elements in the disease category could be considered equivalent provided that the other coordinates of the

facet string (e.g., A, B, and D) were the same. Pulmonary stenosis and coarctation were both low priority diseases. Therefore, an item which requires the student to describe the findings of an electrocardiogram from a patient with pulmonary stenosis (facet string $A_2B_3C_{20}D_3$) was considered equivalent to a question with the same requirements except from a patient with coarctation during the interpretation-of-laboratory-data-phase (Facet string $A_2B_3C_6D_3$). This hypothesis will be tested when the items are used with students.

Due to the criterion of the number of questions per disease, which was not included in the test specification facet strings, the assignment of items to an exam was not straightforward. If a larger pool of items had been developed, it would have been simpler to assign items. Instead, a smaller pool of items was originally developed and the cardiologists were asked to develop specific items once the test development began. However, the cardiologists agreed that developing items from item specification strings was a far simpler and less time consuming task than developing items from objectives without specific guidelines.

The original intent was to develop four parallel tests. Due to logistical problems, and level of difficulty of some items, only two exams were actually used. Although this increases the number of times each item is used, it does not violate the theoretical methodology, nor the notion that the tests are parallel. This revision of the original plan shows the strength and flexibility of facet analysis for item development in that plans can be changed throughout. It would also be possible later to construct additional tests using the same item specification as the original plan.

Recently, Governor Carey of New York signed a "truth-in-testing" bill which permits students to see their graded tests and the correct answers. Obviously, once students can see the actual exams, these items cannot be used again. Test specialists have been able to establish that alternate forms of an exam are

parallel on the basis of consistent performance of different examinees. If items cannot be used repeatedly, then test specialists will need new ways to establish that alternative forms are indeed parallel. The method discussed in this paper is a potentially useful way of maximizing the probability of obtaining parallel forms which can be evaluated later. This facet design analysis might be especially useful for classroom exams.

The use of facet design appears to be a practical way to systematically develop equivalent forms of tests based on a logical analysis of the domain to be tested. This analysis ensures that tests are representative of the learning objectives. Since the actual development of a mapping sentence is not a time consuming activity, it is a viable strategy for content domain specification. If others agree that the mapping sentence logically represents the universe to be tested, then the a priori equivalence of the resulting tests can be established. The development of a large pool of items is a straightforward task when a facet string defines the content of each item. Thus, through the use of facet design, there is a logical justification to the claim that the alternative tests are equivalent in terms of content.

REFERENCES

Berk, R. A. The Application Of Structureal Facet Theory to Achievement Test Construction. Educational Research Quarterly, 1978, 3.

Bloom, B. S. et. al., Taxonomy of Educational Objectives, Handbook I: Cognitive Doman. New York: David McKay, 1956.

Bormuth, J. R. On the Theory of Achievement Test Items. Chicago: University of Chicago Press, 1970.

Engel, J. D. The Development and Investigation of a Methodology to Systematically Construct Multiple Choice Achievement Test Items. Unpublished dissertation at the University of Delaware, 1975.

Gagne, R. W. The Conditions of Learning, 2nd Ed. New York: Holt, Rinehardt, and Winston, 1970.

Guttman, L. Integration of Test Design and Analysis in Proceedings of the 1969 Invitational Conference on Testing Problems. Princeton, N. J.: Educational Testing Service, 1969.

Jordan, J. E. "Facet Theory and the Study of Behavior," in S. Shye, Ed. Theory Construction and Data Analysis in the Behavioral Sciences. San Francisco: Jossey-Bass Publishers, 1978.

Martuza, V. R. Applying Norm-Referenced and Criterion-Referenced Measurement in Education. Boston: Allyn and Bacon, Inc. 1977.

Millman, J. Criterion-Referenced Measurement in W. J. Popham, (Ed). Evaluation in Education. Washington, D.C., American Educational Research Association, 1974.

Popham, W. J. Educational Evaluation. Englewood Cliffs, Prentice-Hall, Inc. 1975.

Shye, S. (Ed.) Theory Construction and Data Analysis in the Behavioral Sciences. San Francisco: Jossey-Bass Publishers, 1978.

Thorndike, Robert L. Educational Measurement. Washington, D.C.: American Council on Education, 1971.

Table 1

Test Specification Table Defined by Facet Strings

Facet String	Number of Questions	Facet String	Number of Questions
A1B1D1	1	A4B1D1	1
A1B1D2	2	A4B1D3	1
A1B1D3	1	A4B1D4	2
A1B1D4	3*	A4B2D1	1
A1B1D5	2	A4B2D3	2
A1B1D6	4*	A4B2D4	1
A1B2D1	1	A4B2D5	1
A1B2D2	1	A4B3D3	1
A1B2D4	2	A4B3D4	1
A1B2D5	1		
A1B2D6	3*	A5B1D1	1
A1B3D1	2	A5B1D2	1
A1B3D2	2	A5B1D5	2
A1B3D4	2	A5B1D4	2
A1B3D5	1*	A5B2D4	2
A1B3D6	3	A5B2D5	2
		A5B3D2	1
A2B1D1	2	A5B3D3	1
A2B1D3	4*	A5B3D4	2
A2B2D1	1	A5B3D5	2
A2B3D3	4*	A5B3D6	2
A2B3D1	1		
A2B3D3	3*	A6B1D5	2
		A6B1D6	1
A3B1D2	2	A6B3D2	1
A3B1D3	1	A6B3D4	1
A3B1D4	1		
A3B2D1	1		
A3B2D2	2		
A3B2D4	1		
A3B2D5	2		
A3B2D6	1		
A3B3D2	1		
A3B3D3	1		
A3B3D4	2		
A3B3D5	2		

The three facets are: A. type of multiple-choice question, B. prioritized objective and D. the step within the clinical process model. The number after each facet letter refers to the specific element of the facet as identified on Figure 2.

*More than two questions were desired for these facet strings because the cardiologists wanted to emphasize interpreting laboratory data, diagnosing diseases and pharmaceutical management regimens generally prescribed. By the nature of these questions they had to be low level types of questions. i.e., either recall information or verbally describe visual information.

Table 2

Exam A Item Specification Table Defined by facet Strings

1. A4C13D1
2. A2C7D1
3. A2C15D1
4. A4C9404
5. A2C11D3
6. A2C13D3
7. A4C20D4
8. A4C3D4
9. A2C3D3
10. A5C15D4
11. A1C1D4
12. A1C2D4
13. A1C3D4
14. A5C24D2
15. A1C7D6
16. A1C2D6
17. A1C14D3
18. A5C15D5
19. A6C18D2
20. A1C2401
21. A6C5D6
22. A3C3D3
23. A6C6D4
24. A1C13D6
25. A1C10D5
26. A1C4D6
27. A6C10D5
28. A1C5D4
29. A6C5D5
30. A1C5D6
31. A5C16D4
32. A1C24D5
33. A4C18D5
34. A1C2401
35. A3C10D2
36. A3C10D2
37. A1C6D4
38. A1C4D4
39. A3C24D5
40. A3C1D2
41. A4C18D4
42. A4C4D4
43. A2C22D3
44. A5C9D2
45. A2C1D3
46. A3C4D4
47. A5C15D4
48. A5C15D4
49. A5C12D4
50. A3C13D4
51. A3C17D4
52. A3C4D5
53. A3C12D2
54. A5C12D5
55. A3C18D1
56. A3C12D6
57. A4C18D1
58. A5C23D3
59. A2C20D3
60. A2C6D3
61. A2C10D3
62. A2C10D3
63. A2C4D3
64. A2C1D3
65. A5C22D6
66. A1C19D2
67. A5C11D5
68. A5C23D5
69. A1C7D5
70. A5C11D4
71. A1C10D2
72. A1C24D2
73. A3C24D2
74. A1C10D4
75. A3C16D5
76. A2C5D1
77. A4C6D3
78. A4C3D3
79. A4C18D3
80. A5C15D4
81. A1C14D1
82. A1C10D6
83. A5C22D6
84. A5C6D5
85. A3C7D4
86. A1C24D5
87. A4C11D3
88. A2C11D3
89. A2C4D3
90. A3C24D5
91. A1C3D6
92. A1C7D6
93. A1C22D2
94. A5C23D1
95. A1C3D1
96. A1C23D6
97. A3C14D4
98. A1C24D6
99. A3C18D2
100. A3C23D3

The three facets are: A. type of multiple choice question, C. name of disease and D. the step within the clinical process model. The number after each facet letter refers to the specicic element of the facet as identified on Figure 2.

TABLE 1

TEST SPECIFICATION TABLE DEFINED BY FACET STRINGS

FACET STRING	NUMBER OF QUESTIONS	FACET STRING	NUMBER OF QUESTIONS
A1B1D1	1	A4B1D1	1
A1B1D2	2	A4B1D3	1
A1B1D3	1	A4B1D4	2
A1B1D4	3*	A4B2D1	1
A1B1D5	2	A4B2D3	2
A1B1D6	4*	A4B2D4	1
A1B2D1	1	A4B2D5	1
A1B2D2	1	A4B3D3	1
A1B2D4	2	A4B3D4	1
A1B2D5	1		
A1B2D6	3*	A5B1D1	1
A1B3D1	2	A5B1D2	1
A1B3D2	2	A5B1D5	2
A1B3D4	2	A5B1D4	2
A1B3D5	1*	A5B2D4	2
A1B3D6	3	A5B2D5	2
		A5B3D2	1
A2B1D1	2	A5B3D3	1
A2B1D3	4*	A5B3D4	2
A2B2D1	1	A5B3D5	2
A2B3D3	4*	A5B3D6	2
A2B3D1	1		
A2B3D3	3*	A6B1D5	2
		A6B1D6	1
A3B1D2	2	A6B3D2	1
A3B1D3	1	A6B3D4	1
A3B1D4	1		
A3B2D1	1		
A3B2D2	2		
A3B2D4	1		
A3B2D5	2		
A3B2D6	1		
A3B3D2	1		
A3B3D3	1		
A3B3D4	2		
A3B3D5	2		

The three facets are: A. type of multiple-choice question. B. prioritized objectives and C. the step within the clinical process model. The number after each facet letter refers to the specific element of the facet as identified on Figure 2.

More than two questions were desired for these facet strings because the cardiologists wanted to emphasize interpreting laboratory data, diagnosing diseases and pharmaceutical management regimens generally prescribed. By the nature of these questions they had to be low level types of questions. i.e., either recall information or verbally described visual information.

TABLE 2

Exam A Item Specification Table Defined by Facet Strings

1. A4C13D1
2. A2C7D1
3. A2C15D1
4. A4C9A04
5. A2C11D3
6. A2C13D3
7. A4C20D4
8. A4C3D4
9. A2C3D3
10. A5C15D4
11. A1C1D4
12. A1C2D4
13. A1C3D4
14. A5C24D2
15. A1C7D6
16. A1C2D6
17. A1C14D3
18. A5C15D5
19. A6C18D2
20. A1C24D1
21. A6C5D6
22. A3C3D3
23. A6C6D4
24. A1C13D6
25. A1C10D5
26. A1C4D6
27. A6C10D5
28. A1C5D4
29. A6C5D5
30. A1C5D6
31. A5C16D4
32. A1C24D5
33. A4C18D5
34. A1C24D1
35. A3C10D2
36. A3C10D2
37. A1C6D4
38. A1C4D4
39. A3C24D5
40. A3C1D2
41. A4C18D4
42. A4C4D4
43. A2C22D3
44. A5C9D2
45. A2C1D3
46. A3C4D4
47. A5C15D4
48. A5C15D4
49. A5C12D4
50. A3C13D4
51. A3C17D4
52. A3C4D5
53. A3C12D2
54. A5C12D5
55. A3C18D1
56. A3C12D6
57. A4C18D1
58. A5C23D3
59. A2C20D3
60. A2C6D3
61. A2C10D3
62. A2C10D3
63. A2C4D3
64. A2C1D3
65. A5C22D6
66. A1C19D2
67. A5C11D5
68. A5C23D5
69. A1C7D5
70. A5C11D4
71. A1C10D2
72. A1C24D2
73. A3C24D2
74. A1C10D4
75. A3C16D5
76. A2C5D1
77. A4C6D3
78. A4C3D3
79. A4C18D3
80. A5C15D4
81. A1C14D1
82. A1C10D6
83. A5C22D6
84. A5C6D5
85. A3C7D4
86. A1C24D5
87. A4C11D3
88. A2C11D3
89. A2C4D3
90. A3C24D5
91. A1C3D6
92. A1C7D6
93. A1C22D2
94. A5C23D1
95. A1C3D1
96. A1C23D6
97. A3C14D4
98. A1C24D6
99. A3C18D2
100. A3C23D3

The three facets are:. A. type of multiple choice question, C. name of disease and D. the step within the clinical process model. The number after each facet letter refers to the specific element of the facet as identified on Figure 2.

FIGURE 1

A SAMPLE MAPPING SENTENCE (Millman, 1974)

A. REFERENT

1. self
2. music education
3. general public
4. students

B. REFERENTS COMMITMENT

1. belief
2. feeling

The subject contributes to the [A] the [B]

C. LEVEL

1. elementary
2. secondary

D. ACTORS (e.g. music curriculum's) BEHAVIOR

1. should be
2. is

that the [C] school music curriculm [D] taught by

E. TEACHER

1. classroom
2. specialist

F. GENERAL STUDENT NEEDS

1. relaxation
2. means of expression
3. breaks from academics
4. emotional stimulation
5. self-discipline
6. fun time
7. contact with a human
8. group activity
9. uncover unknown talents
10. public performance
11. creative outlet
12. success
13. bring out shy students

[E] for the purpose of fulfilling [F]

---

1. A through F are facets.

2. A1 through A4 are facet elements.

3. A1, B1, C1, D2, E2, F3 is a facet string.

4. An item based on the facet string in #3 above is:

   Do you believe that the elementary school music curriculum is taught by specialists for the purpase of giving students a break from academics?

Figure 2

In answering multiple choice items on a cardiology exam, students will

A. Employ cognitive Process

- A_1 recall factual information
- A_2 verbally describe visual information
- A_3 interpret factual information
- A_4 interpret visual information
- A_5 interpret information from various sources
- A_6 apply information from various sources

when confronted with a

B. Disease priority category

- B_1 high priority
- B_2 medium priority
- B_3 low priorty

which presents as

C. Specific Disease

- C_1 acute MR
- C_2 AR
- C_3 AS
- C_4 ASD
- C_5 Angina - CAD
- C_6 Coarctation
- C_7 Congestive Cardiomyopathy
- C_8 Endocarditis
- C_9 Functional M
- C_{10} Hypertension
- C_{11} IIISS
- C_{12} MVP
- C_{13} MR
- C_{14} MS
- C_{15} Normal patient
- C_{16} PDA
- C_{17} Pericarditis
- C_{18} Pul Hypertension-primary
- C_{19} Pulm. Emboli
- C_{20} Pulmonary Stenosis
- C_{21} Rheumatic fever
- C_{22} Vent Aneurysm
- C_{23} VSD

at the stage when

D. The step involved within the clinical process

- D_1 data gathering
- D_2 order laboratory data
- D_3 interpret laboratory data
- D_4 diagnose patients problems
- D_5 determine management plan
- D_6 monitor on-going management

is required

Test specification involves Facets A, C, and D.
Item specification involves Facets A, B, and D.